

AD-A231 338

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188REPORT SECURITY CLASSIFICATION  
UNCLASSIFIED1b RESTRICTIVE MARKINGS  
SELECTED

SECURITY CLASSIFICATION AUTHORITY

JAN 22 1991

DECLASSIFICATION/DOWNGRADING SCHEDULE

3 DISTRIBUTION/AVAILABILITY OF REPORT  
This document has been approved for public release and sale; its distribution is unlimited.

PERFORMING ORGANIZATION REPORT NUMBER(S)

IBM Research Report RJ 7794

5. MONITORING ORGANIZATION REPORT NUMBER(S)

- Technical Report No. 27

NAME OF PERFORMING ORGANIZATION

IBM Research Division  
Almaden Research Center6b OFFICE SYMBOL  
(if applicable)

ADDRESS (City, State, and ZIP Code)

650 Harry Road  
San Jose, CA 95120-6099

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

7b. ADDRESS (City, State, and ZIP Code)

Chemistry Division Code 1113  
Arlington, VA 22217

8a. NAME OF FUNDING/SPONSORING ORGANIZATION

Office of Naval Research

8b. OFFICE SYMBOL  
(if applicable)

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-84-C-0708, 4131022

8c. ADDRESS (City, State, and ZIP Code)

Chemistry Division, Code 1113  
Arlington, VA 22217

10. SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NOPROJECT  
NOTASK  
NOWORK UNIT  
ACCESSION NO

11. TITLE (Include Security Classification)

Observation of Spectral Diffusion in Crystals Using Single Impurity Molecules

12. PERSONAL AUTHOR(S)

W. P. Ambrose and W. E. Moerner

13a. TYPE OF REPORT

interim technical

13b. TIME COVERED

FROM TO

14. DATE OF REPORT (Year, Month, Day)

1990, 10, 31

15. PAGE COUNT

13

16. SUPPLEMENTARY NOTATION

17

COSATI CODES

FIELD

GROUP

SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

single center detection and spectroscopy  
spectral diffusion, hole-burning

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Early measurements of the optical absorption spectrum of a single impurity molecule in a solid at low temperatures, while intriguing, nevertheless produced insufficient signal-to-noise ratio for the observation of new physical effects. Recent advances in high-efficiency fluorescence excitation spectroscopy have removed this obstacle. For the case of pentacene molecules in p-terphenyl, both stable as well as spectrally diffusing single molecules can be observed.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Dr. W.E. Moerner

22b. TELEPHONE (Include Area Code)

(408) 927-2426

22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete  
S/N 0102-LF-014-6603SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED

91 1 22 062

OFFICE OF NAVAL RESEARCH

Contract N00014-84-C-0708

R&T Code 4131022

Technical Report No. 27

**Observation of Spectral Diffusion in Crystals Using Single  
Impurity Molecules**

by

W. P. Ambrose and W. E. Moerner

Prepared for Publication

in

Nature

IBM Research Division  
Almaden Research Center  
650 Harry Road  
San Jose, California 95120-6099

October 31, 1990

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

RJ 7794 (71911) 10/31/90  
Physics

## OBSERVATION OF SPECTRAL DIFFUSION IN CRYSTALS USING SINGLE IMPURITY MOLECULES

W. P. Ambrose and W. E. Moerner

IBM Research Division  
Almaden Research Center  
San Jose, California 95120

Early measurements of the optical absorption spectrum of a single impurity molecule in a solid at low temperatures<sup>1</sup>, while intriguing, nevertheless produced insufficient signal-to-noise ratio (S/N) for the observation of new physical effects. Recent advances in high-efficiency fluorescence excitation spectroscopy<sup>2</sup> with ultrathin sublimed crystals have removed this obstacle. For the case of pentacene impurities in crystals of *p*-terphenyl, we observe two radically different behaviors. For class I guest molecules, the center frequency does not change with time and the low-power linewidth at 1.5 K is in agreement with previous work<sup>3</sup>. However, for class II molecules which are only found far out in the wings of the inhomogeneous line, the center frequency spontaneously jumps from one value to several other possible values on a time scale ranging from seconds to minutes, independent of laser power. This surprising effect is analogous to spectral diffusion processes that play a crucial role in the physics of the amorphous state<sup>4,5</sup>; here, the single-molecule technique allows the spectral changes to be followed in real time.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Early measurements of the optical absorption spectrum of a single impurity molecule in a solid at low temperatures<sup>1</sup>, while intriguing, nevertheless produced insufficient signal-to-noise ratio (S/N) for the observation of new physical effects. Recent advances in high-efficiency fluorescence excitation spectroscopy<sup>2</sup> with ultrathin sublimed crystals have removed this obstacle. For the case of pentacene impurities in crystals of *p*-terphenyl, we observe two radically different behaviors. For class I guest molecules, the center frequency does not change with time and the low-power linewidth at 1.5 K is in agreement with previous work<sup>3</sup>. However, for class II molecules which are only found far out in the wings of the inhomogeneous line, the center frequency spontaneously jumps from one value to several other possible values on a time scale ranging from seconds to minutes, independent of laser power. This surprising effect is analogous to spectral diffusion processes that play a crucial role in the physics of the amorphous state<sup>4,5</sup>: here, the single-molecule technique allows the spectral changes to be followed in real time.

In our fluorescence excitation technique, a tunable dye laser (1 MHz linewidth) was used to excite the 0-0 electronic transition of pentacene substitutional impurities in the *p*-terphenyl crystal. As the excitation wavelength was varied, the emitted fluorescence that passed through a long pass filter was measured with a photon counter. While such methods are capable of measuring the extremely small absorptions from single ions in traps<sup>6</sup>, in a single-molecule experiment in a solid the scattered light from the sample and other components must be reduced to very low levels. This was achieved by limiting the excited volume with 1-10  $\mu\text{m}$  thick sublimed single crystal samples and laser focal diameters of 5  $\mu\text{m}$  full width at half-maximum (FWHM). Following Orrit et al.<sup>2</sup>, the Stokes-shifted fluorescence was collected with high efficiency using 0.98 numerical aperture optics.

The crystals were cooled to 1.5 K where the homogeneous linewidth of individual pentacene defects for the O<sub>1</sub> site<sup>7</sup> ( $\gamma \approx 7.5 \text{ MHz}$ <sup>3</sup>) is much less than the inhomogeneous line

width ( $\Gamma_1 \approx 2-42$  GHz, depending upon strain in the crystal). This well-known inhomogeneous broadening effect<sup>9</sup> occurs for all zero-phonon transitions of defects in solids at low temperatures<sup>10</sup>. The inhomogeneous line profile represents the center frequency distribution of the pentacene guests produced by random strain and electric fields present in the crystal. For single-molecule spectra, the laser may be tuned far into the wings of the inhomogeneous line where the number of defects per homogeneous width is less than one; i.e., where non-overlapping individual pentacene defects may be found. For most measurements, the pentacene concentration was approximately  $6.6 \times 10^{-4}$  moles/mole, and  $\Gamma_1 \approx 6$  GHz. To record single-molecule spectra we used laser wavelengths  $\approx 0.2$  nm from the inhomogeneous line center for the O<sub>1</sub> 0-0 site origin at 592.321 nm<sup>7</sup>.

Fig 1. shows excitation spectra taken at a series of laser focal spot positions. The three largest peaks isolated in both position and frequency are single pentacene defects of class I. The width of the peaks along the position axis (FWHM = 5  $\mu$ m) is a direct measure of the intensity profile of the focused laser beam in the crystal. The width of the peaks in frequency varies from 8 to 13 MHz as the laser intensity at the defect varies with position. This broadening is due to saturation of the defect optical transition. Using three-level saturation expressions<sup>11</sup>, the saturation intensity of these defects is estimated to be  $1.8 \pm 0.8$  mW/cm<sup>2</sup>, which is a factor of 40 lower than the 70 mW/cm<sup>2</sup> value one would calculate from pentacene photophysical parameters including intersystem crossing<sup>12</sup>. Apparently (and not surprisingly), the local pentacene environment this far out in the wings of the line produces alterations in the intersystem crossing rates. In addition to the class I peaks, there are smaller features in Fig. 1 that do not follow a definite trend with position suggesting a time dependence from scan to scan.

Fig. 2 shows a typical time sequence of excitation spectra obtained at a fixed laser spot position. The spectra contain three time-independent class I peaks, which form stable ridges

along the time axis near -130, -70, and +10 MHz. In the 0 to 150 MHz region is a single class II defect whose frequency varies from one scan to the next. Indeed, this same molecule can appear several times in the same 2 min laser scan, or almost not at all. Notice that in the same  $\approx 50 \mu\text{m}^3$  volume of crystal and at nearly the same frequency there are both class I defects as well as spectrally migrating class II defects.

If the laser frequency in Fig. 2 is fixed at any detuning from 0 to +100 MHz, the collected fluorescence turns on and off discontinuously like a random telegraph<sup>13</sup> or quantum jump<sup>14</sup> signal. However, the absence of fluorescence only means that the defect is not absorbing at the laser frequency. To follow the resonance frequency directly, we scanned more rapidly and sacrificed S/N in order to measure the spectral position of the defect once each second. Fig. 3 (upper part) shows how the resonance frequency of a spectrally jumping class II defect evolves with time. A computer locates the peak in each scan and records this resonance frequency as a function of time as a trend in the lower part of Fig. 3. The optical transition energy appears to have a preferred set of values and performs spectral jumps between these values that are discontinuous on the 1 sec time scale of the measurement.

Using several samples from the same sublimation run, a variety of class II defects have been observed that have two or more preferred frequencies spaced by 20 to 60 MHz with mean times between jumps of 1 to 420 seconds. Moreover, the occurrence of class II behavior is quite common in the wings of the inhomogeneous line (to the red as well as to the blue). By observing 133 single molecules at several positions to the red of  $O_1$ , the fraction of class II increases from 13% at 592.365 nm to 40% at 592.550 nm. On the other hand, using a thin, strain-free, lightly doped sample with concentration  $5 \times 10^{-9}$  moles/mole and  $\Gamma_1 = 2$  GHz, single molecules could be observed 0.006 nm from line center and only class I behavior was found. We conclude that class II behavior is related to the improbable, highly strained local environments that occur far out in the wings of the inhomogeneous line.

The mean time between spectral jumps varies no more than 20% for laser intensities from 3.3 to 65 mW/cm<sup>2</sup>. Therefore, the spontaneous spectral jumps do not appear to be photo-induced changes such as that which occurs when a persistent spectral hole is formed near the center of the inhomogeneous band<sup>15</sup>. By analogy to glasses, we call the spectral jumping effect spectral diffusion, because the absorbing molecule appears to be coupled to an ensemble of diagonal (energy-changing) perturbations that shift the resonance frequency in a stochastic manner. On the other hand, class I defects have not been observed to jump with the laser frequency fixed at the resonance frequency for periods up to 10 min, or by scanning and detecting the resonance at regular intervals up to 55 min. Thus we identify class I defects as isolated pentacene molecules that have stable surroundings.

Measurements of the temperature dependence and the power broadening behavior for class I and class II molecules show that both the saturation intensity and dephasing rates differ from center to center and from class to class. However, all centers appear to have the same lifetime-limited  $\gamma \approx 7.8 \pm 0.2$  MHz, in agreement with coherent transient data<sup>3</sup>. By comparison with the measured properties of dimers of pentacene in *p*-terphenyl<sup>8</sup>, we find no strong evidence that either class of defect represents dimers. Individual defects at nearly the same frequency far in the wings of the inhomogeneous line each have unique spectral jumping behavior, which suggests different nearby host configurations. We are thus led to propose that class II defects are located in regions of the crystal where a variety of local structural configurations are energetically accessible at low temperatures. These local configurations may be modeled by an ensemble of two-level systems (TLS) in analogy with the glass case<sup>5</sup>. The TLS undergo phonon-assisted tunnelling<sup>16</sup> and the pentacene resonance shifts with changes in the quantum numbers describing the nearby host crystal configuration. One possible source for the TLS would be tipping of the central phenyl ring of *p*-terphenyl among various allowed orientations made available by disorder in the crystal structure<sup>17</sup>. The future proof of this model requires detailed temperature dependent measurements of the jump rate

for a single center over a wide range of temperature in hope of observing the characteristic  $\coth(E/2k_B T)$  dependence, where  $E$  is the energy splitting in an asymmetric TLS model and  $k_B$  is Boltzmann's constant.

With the ability to measure individual defect spectra at 1 sec intervals with reasonable S/N, we find that single pentacene molecules far from the center of the inhomogeneous line in crystals of p-terphenyl exhibit spontaneous spectral jumps at liquid-helium temperatures. Since the spectral jump rates and sizes are different for different defects, direct observation of this spectral diffusion effect would not be possible in any spectroscopy that averages over many defects. Direct measurement of the behavior of individual guest molecules in a glass would lead to a clearer understanding of the origins of amorphous solid dynamics<sup>4, 18</sup> than is available from ensemble-averaged measurements. Much more intriguing is the fact that spectral jumping has been first directly measured in a crystal where such an effect is not expected.



## References

1. W. E. Moerner and L. Kador, *Phys. Rev. Lett.* **62**, 2535-2538 (1989), L. Kador, D. E. Horne, and W. E. Moerner, *J. Phys. Chem.* **94**, 1237-1248 (1990).
2. M. Orrit and J. Bernard, *Phys. Rev. Lett.* (submitted).
3. H. de Vries and D. A. Wiersma, *J. Chem. Phys.* **69**, 897-901 (1978).
4. J. Friedrich and D. Haarer, "Structural Relaxation Processes in Polymers and Glasses as Studied by High Resolution Optical Spectroscopy," in *Optical Spectroscopy of Glasses*, I. Zschokke, Ed. (D. Reidel, 1986), pp. 149-198.
5. See *Amorphous Solids: Low Temperature Properties*, W. A. Phillips, editor (Springer, Berlin, 1981).
6. W. M. Itano, J. C. Bergquist, and D. J. Wineland, *Science* **237**, 612-617 (1987).
7. There are four major sites corresponding to pentacene substitution for each of the four *p*-terphenyl molecules in the unit cell. See R. W. Olson and M. D. Fayer, *J. Phys. Chem.* **84**, 2001-2004 (1980).
8. F. G. Patterson, H. W. H. Lee, W. L. Wilson, and M. D. Fayer, *Chem. Phys.* **84**, 51-60 (1984).
9. A. M. Stoneham, *Rev. Mod. Phys.* **41**, 82-108 (1969).
10. W. M. Yen and P. M. Selzer, eds. *Laser Spectroscopy of Solids*, (Springer, Berlin, Heidelberg, 1981).
11. H. de Vries and D. A. Wiersma, *J. Chem. Phys.* **72**, 1851-1863 (1980).
12. H. de Vries and D. A. Wiersma, *J. Chem. Phys.* **70**, 5807-5822 (1979).
13. C. W. Gardiner, *Handbook of Stochastic Methods* (Springer, Berlin, 1983) pp. 78-79.
14. W. Nagourney, J. Sandberg, and H. Dehmelt, *Phys. Rev. Lett.* **56**, 2797-2799 (1986).
15. W. E. Moerner and T. P. Carter, *Phys. Rev. Lett.* **59**, 2705-2708 (1987).
16. J. A. Sussman, *Phys. kondens. Materie* **2**, 146-160 (1964).

17. J. L. Baudour, Y. Delugeard, and H. Cailleau, *Acta Cryst.* B32, 150-154 (1976).
18. B. Golding and J. E. Graebner, in Ref. 5, pp. 107-134.

### **Acknowledgements**

We thank Prof. Orrit for providing a copy of his manuscript prior to publication. This work was supported in part by the U. S. Office of Naval Research.

## Figure Legends

Figure 1. Single pentacene defect fluorescence excitation spectral landscape in a crystal of p-terphenyl. 0 MHz detuning  $\equiv$  592.544 nm,  $T = 1.5$  K, laser power = 1.5 nW, 2 min. per spectrum.

Figure 2. Single defect fluorescence excitation spectra time dependence. Upper half: wire plot of individual spectra, lower half: contour plot of same data, 2 min. per spectrum, laser power = 1.5 nW, and 0 MHz detuning  $\equiv$  592.539 nm. The appearance of class I at negative detunings and class II at positive detunings has no significance.

Figure 3. Spectral jumps in the resonance frequency of a class II single pentacene defect. Upper half: individual 1 sec. spectra, lower half: time dependence of center frequency, 0 MHz detuning  $\equiv$  592.546 nm.




Figure 1




Figure 2




Figure 3

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)  
Chemistry Division, Code 1113  
800 North Quincy Street  
Arlington, Virginia 22217-5000

Commanding Officer (1)  
Naval Weapons Support Center  
Dr. Bernard E. Douda  
Crane, Indiana 47522-5050

Dr. Richard W. Drisko (1)  
Naval Civil Engineering  
Laboratory  
Code L52  
Port Hueneme, CA 93043

David Taylor Research Center (1)  
Dr. Eugene C. Fischer  
Annapolis, MD 21402-5067

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)  
Chemistry Division, Code 385  
Naval Weapons Center  
China Lake, CA 93555-6001

Chief of Naval Research (1)  
Special Assistant for Marine  
Corps Matters  
Code 00MC  
800 North Quincy Street  
Arlington, VA 22217-5000

Dr. Bernadette Eichinger (1)  
Naval Ship Systems Engineering  
Station  
Code 053  
Philadelphia Naval Base  
Philadelphia, PA 19112

Dr. Sachio Yamamoto (1)  
Naval Ocean Systems Center  
Code 52  
San Diego, CA 92152-5000

Dr. Harold H. Singerman (1)  
David Taylor Research Center  
Code 283  
Annapolis, MD 21402-5067

+ DTIC (2)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

ENCLOSURE(-)